

Connecting via Winsock to STN

10/687,924h

Welcome to STN International! Enter x:x

structure search

$G_1 = 0,5$   
 $G_2 = C, N$

LOGINID:SSSPTAAJP1626

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2



\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 FEB 25 CA/CAPLUS - Russian Agency for Patents and Trademarks  
(ROSPATENT) added to list of core patent offices covered  
NEWS 4 FEB 28 PATDPAFULL - New display fields provide for legal status  
data from INPADOC  
NEWS 5 FEB 28 BABS - Current-awareness alerts (SDIs) available  
NEWS 6 FEB 28 MEDLINE/LMEDLINE reloaded  
NEWS 7 MAR 02 GBFULL: New full-text patent database on STN  
NEWS 8 MAR 03 REGISTRY/ZREGISTRY - Sequence annotations enhanced  
NEWS 9 MAR 03 MEDLINE file segment of TOXCENTER reloaded  
NEWS 10 MAR 22 KOREAPAT now updated monthly; patent information enhanced  
NEWS 11 MAR 22 Original IDE display format returns to REGISTRY/ZREGISTRY  
NEWS 12 MAR 22 PATDPASPC - New patent database available  
NEWS 13 MAR 22 REGISTRY/ZREGISTRY enhanced with experimental property tags  
NEWS 14 APR 04 EPFULL enhanced with additional patent information and new  
fields  
NEWS 15 APR 04 EMBASE - Database reloaded and enhanced

NEWS EXPRESS JANUARY 10 CURRENT WINDOWS VERSION IS V7.01a, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 10 JANUARY 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 15:03:24 ON 06 APR 2005

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 15:03:33 ON 06 APR 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1  
DICTIONARY FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.


```
*****
*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*
*****
```

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10681924h.str



chain nodes :

11 15

ring nodes :

1 2 3 4 5 6 7 8 9

chain bonds :

6-15 8-11

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-9 7-8 8-9

exact/norm bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-9 6-15 7-8 8-9 8-11

G1:O,S

G2:C,N

G3:cy,Ak

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 11:CLASS  
15:CLASS

L1           STRUCTURE UPLOADED

=> s L1

SAMPLE SEARCH INITIATED 15:03:47 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 11643 TO ITERATE

8.6% PROCESSED       1000 ITERATIONS                           0 ANSWERS  
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)  
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS:   ONLINE   \*\*COMPLETE\*\*  
                          BATCH    \*\*COMPLETE\*\*  
PROJECTED ITERATIONS:       226395 TO   239325  
PROJECTED ANSWERS:           0 TO           0

L2           0 SEA SSS SAM L1

=> s L1 full

FULL SEARCH INITIATED 15:03:52 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 233513 TO ITERATE

100.0% PROCESSED   233513 ITERATIONS                       32 ANSWERS  
SEARCH TIME: 00.00.06

L3           32 SEA SSS FUL L1

=> fil caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	161.33	161.54

FILE 'CAPLUS' ENTERED AT 15:04:03 ON 06 APR 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 6 Apr 2005   VOL 142 ISS 15  
FILE LAST UPDATED: 5 Apr 2005   (20050405/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s L3

L4

7 L3

=&gt; d 1-7 ibib abs hitstr

L4 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:331928 CAPLUS

DOCUMENT NUMBER: 140:357354

TITLE: A preparation of benzimidazolone derivatives useful as anti-inflammatory agents

INVENTOR(S): Dhar, T. G. Murali; Potin, Dominique; Maillet, Magali  
Jeannine Blandine; Launay, Michele; Nicolai, Eric  
Antoine; Iwanovicz, Edwin J.

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 69 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004032861	A2	20040422	WO 2003-US31960	20031009
WO 2004032861	A3	20040805		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2004116467	A1	20040617	US 2003-681924	20031009
PRIORITY APPLN. INFO.:			US 2002-417935P	P 20021011
OTHER SOURCE(S):	MARPAT 140:357354			
GI				

our  
app.

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB The invention relates to benzimidazolone derivs. of formula I [wherein: K is O or S; Q is a bond or C(O), etc.; Ar is (un)substituted (hetero)aryl; J1 is a bond, -N(R4)-, etc.; J2 and J3 are -N(R4)- or (un)substituted CH2, etc.; Y and Z are independently selected from N, (un)substituted CH, etc.; R1 = H, (un)substituted alk(en)yl, (hetero)aryl, cycloalkyl, etc.; R2 and R3 are independently selected from H, halogen, NO2, CN, (un)substituted alk(en)yl, etc.; R4 is H, (un)substituted alk(en)yl, CN, C(O)-alkyl, O-alkyl, etc.], their enantiomers, diastereomers, and pharmaceutically-acceptable salts, useful as anti-inflammatory agents. Compds. I were tested in an H1-HeLa adhesion assay and in a HUVEC (human umbilical vein endothelial cells) adhesion assay (no biol. data). For instance, benzimidazole derivative II was prepared via intramol. heterocyclization of the obtained urea derivative III, and N-acetylation of the obtained benzimidazole derivative IV (no yield data).


IT 681261-14-3P 681261-15-4P 681261-21-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(intermediate; preparation of benzimidazolone derivs. useful as anti-inflammatory agents)


RN 681261-14-3 CAPLUS

CN 2H-Benzimidazol-2-one, 3a-[(4-bromophenyl)methyl]-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro- (9CI) (CA INDEX NAME)




RN 681261-15-4 CAPLUS

CN Benzonitrile, 4-[[1-(3,5-dichlorophenyl)-1,2,3,4,5,6-hexahydro-2-oxo-3aH-benzimidazol-3a-yl]methyl]- (9CI) (CA INDEX NAME)



RN 681261-21-2 CAPLUS

CN 1H-Benzimidazole-1-acetic acid, 7a-[(4-cyanophenyl)methyl]-3-(3,5-dichlorophenyl)-2,3,5,6,7,7a-hexahydro-2-oxo-, ethyl ester (9CI) (CA INDEX NAME)



IT 681261-16-5P 681261-17-6P 681261-18-7P  
681261-19-8P 681261-20-1P 681261-22-3P


681261-23-4P 681261-24-5P 681261-25-6P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of benzimidazolone derivs. useful as anti-inflammatory agents)


RN 681261-16-5 CAPLUS

CN Benzonitrile, 4-[[1-(3,5-dichlorophenyl)-1,2,3,4,5,6-hexahydro-3-methyl-2-oxo-3aH-benzimidazol-3a-yl]methyl]- (9CI) (CA INDEX NAME)




RN 681261-17-6 CAPLUS

CN 2H-Benzimidazol-2-one, 3a-[(4-bromophenyl)methyl]-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-3-methyl- (9CI) (CA INDEX NAME)




RN 681261-18-7 CAPLUS

CN 2H-Benzimidazol-2-one, 3-acetyl-3a-[(4-cyanophenyl)methyl]-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro- (9CI) (CA INDEX NAME)




RN 681261-19-8 CAPLUS

CN 2H-Benzimidazol-2-one, 3-acetyl-3a-[(4-bromophenyl)methyl]-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro- (9CI) (CA INDEX NAME)




RN 681261-20-1 CAPLUS

CN Benzonitrile, 4-[[1-(3,5-dichlorophenyl)-3-ethyl-1,2,3,4,5,6-hexahydro-2-oxo-3aH-benzimidazol-3a-yl]methyl]- (9CI) (CA INDEX NAME)




RN 681261-22-3 CAPLUS

CN 1H-Benzimidazole-1-acetic acid, 7a-[(4-cyanophenyl)methyl]-3-(3,5-dichlorophenyl)-2,3,5,6,7,7a-hexahydro-2-oxo- (9CI) (CA INDEX NAME)




RN 681261-23-4 CAPLUS

CN 1H-Benzimidazole-1-acetic acid, 7a-[(4-bromophenyl)methyl]-3-(3,5-dichlorophenyl)-2,3,5,6,7,7a-hexahydro-2-oxo-, ethyl ester (9CI) (CA INDEX NAME)



RN 681261-24-5 CAPLUS

CN 1H-Benzimidazole-1-hexanoic acid, 7a-[(4-cyanophenyl)methyl]-3-(3,5-dichlorophenyl)-2,3,5,6,7,7a-hexahydro-2-oxo-, ethyl ester (9CI) (CA INDEX NAME)




RN 681261-25-6 CAPLUS




CN1C(=O)N(C2=CC(=CC(=C2)Cl)Cl)C3=CC=C(C=C3C4=CC=CC=C4C5=NC=CC=N5)C6=CC=CC=C6

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxamide, 2,5,6,7-tetrahydro-N,3,5,7-tetramethyl-6-oxo- (9CI) (CA INDEX NAME)




RN 181585-51-3 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxamide, N-ethyl-2,5,6,7-tetrahydro-3,5,7-trimethyl-6-oxo- (9CI) (CA INDEX NAME)




RN 181585-56-8 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-3,4a-dicarboxamide, 2,5,6,7-tetrahydro-N,N',5,7-tetramethyl-6-oxo- (9CI) (CA INDEX NAME)




RN 181585-57-9 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-3,4a-dicarboxamide, N,N'-diethyl-2,5,6,7-tetrahydro-5,7-dimethyl-6-oxo- (9CI) (CA INDEX NAME)




RN 181585-58-0 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-3,4a-dicarboxamide, 2,5,6,7-tetrahydro-5,7-dimethyl-6-oxo-N,N'-dipropyl- (9CI) (CA INDEX NAME)




RN 181585-59-1 CAPLUS  
 CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-3,4a-dicarboxamide, 2,5,6,7-tetrahydro-5,7-dimethyl-N,N'-bis(2-methylpropyl)-6-oxo- (9CI) (CA INDEX NAME)




L4 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1989:430150 CAPLUS  
 DOCUMENT NUMBER: 111:30150  
 TITLE: Isolation and characterization of 5-[3'-(7',9'-dimethyluric acid)]-7,9-dimethyl-Δ3,4-isouric acid. An unstable product of electrochemical oxidation of 7,9-dimethyluric acid  
 AUTHOR(S): Subramanian, P.; Dryhurst, Glenn  
 CORPORATE SOURCE: Dep. Chem. Biochem., Univ. Oklahoma, Norman, OK, 73019, USA  
 SOURCE: Journal of Electroanalytical Chemistry and Interfacial Electrochemistry (1989), 262(1-2), 281-7  
 CODEN: JEIEBC; ISSN: 0022-0728  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB An account is given of the isolation and structural characterization of the title compound (I) as well as mechanisms for its formation and decomposition  
 The starting material, 7,9-dimethyluric acid (II), was of com. origin and the electrochem. apparatus was that described earlier. Controlled potential electrooxidn. of II was carried out at 0.65 V in NH<sub>4</sub>OAc as the supporting electrolyte. To assay the results HPLC, electron impact mass spectrometry, fast-atom bombardment mass spectrometry and similar methods were used.  
 IT 121263-20-5P 121350-86-5P  
 RL: RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
 (synthesis of, mechanisms of formation and decomposition in relation to)  
 RN 121263-20-5 CAPLUS  
 CN 1H-Purine-2,6,8(3H)-trione, 3-(1,2,6,7,8,9-hexahydro-7,9-dimethyl-2,6,8-

trioxo-5H-purin-5-yl)-7,9-dihydro-7,9-dimethyl-, stereoisomer (9CI) (CA INDEX NAME)



RN 121350-86-5 CAPLUS

CN 1H-Purine-2,6,8(3H)-trione, 3-(1,2,6,7,8,9-hexahydro-7,9-dimethyl-2,6,8-trioxo-5H-purin-5-yl)-7,9-dihydro-7,9-dimethyl-, stereoisomer (9CI) (CA INDEX NAME)



L4 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1988:167430 CAPLUS

DOCUMENT NUMBER: 108:167430

TITLE: Intermediates in the transformation of  
pyrimido[5,4-e]-1,2,4-triazinediones into  
imidazo[4,5-e]-1,2,4-triazinones

AUTHOR(S): Shorshnev, S. V.; Esipov, S. E.; Yakushkina, N. I.;  
Klyuev, N. A.; Zhil'nikov, V. G.; Chernyshev, A. I.

CORPORATE SOURCE: Vses. Nauchno-Issled. Inst. Antibiot., Moscow, 113105,  
USSR

SOURCE: Khimiya Geterotsiklicheskikh Soedinenii (1987), (9),  
1252-9

CODEN: KGSSAQ; ISSN: 0453-8234





DOCUMENT TYPE: Journal

LANGUAGE: Russian

OTHER SOURCE(S): CASREACT 108:167430

GI

*needs to be phys. exptl. Grastelli - ny. 10m ch. 1987*




AB Treating pyrimidotriazinediones I (R = H, Me) with aqueous NaOH gave imidazopyrimidinecarboxylic acids II which were decarboxylated by acid to imidazotriazinones III followed by KMnO<sub>4</sub> dehydrogenation to give imidazotriazinones IV.

IT **97943-27-6P**

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and decarboxylation of)

RN 97943-27-6 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 1,2,5,6-tetrahydro-5-methyl-6-oxo- (9CI) (CA INDEX NAME)




IT **113947-23-2P**

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 113947-23-2 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 2,5,6,7-tetrahydro-5,7-dimethyl-6-oxo-, methyl ester (9CI) (CA INDEX NAME)




IT **97943-30-1P**

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation, decarboxylation and esterification of)


RN 97943-30-1 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 2,5,6,7-tetrahydro-

5,7-dimethyl-6-oxo- (9CI) (CA INDEX NAME)




L4 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1987:156125 CAPLUS  
 DOCUMENT NUMBER: 106:156125  
 TITLE: NMR spectroscopic investigation of reumycin alkaline hydrolysis  
 AUTHOR(S): Shorshnev, S. V.; Esipov, S. E.; Chernyshev, A. I.  
 CORPORATE SOURCE: All-Union Res. Inst. Antibiot., Moscow, USSR  
 SOURCE: Antibiotiki i Meditsinskaya Biotekhnologiya (1987), 32(2), 116-20  
 CODEN: AMBIEH; ISSN: 0233-7525  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 GI




AB The transformation sequence of reumycin (I) in aqueous (D2O) solns. with various pH was studied by NMR spectroscopy and the structures of the products were determined. Formation of 6-(3-methylureido)-1,2,4-triazine-5-carboxylic acid was the 1st stage of I alkaline hydrolysis. The subsequent cyclization of this compound resulted in the formation of 5-methyl-4,4a-dihydro-5H-imidazo[4,5-e]-1,2,4-triazin-6(7H)-one-4a-carboxylic acid mono-, di- or trianionic forms depending on the medium pH and due to dissociation of the carboxylic group, N(4)H group of the triazine ring, and N(7)H group of the imidazolidinone ring.


IT **97943-27-6P**  
 RL: FORM (Formation, nonpreparative); PREP (Preparation)  
 (formation of, in reumycin alkaline hydrolysis)  
 RN 97943-27-6 CAPLUS  
 CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 1,2,5,6-tetrahydro-5-methyl-6-oxo- (9CI) (CA INDEX NAME)




L4 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1985:504756 CAPLUS  
 DOCUMENT NUMBER: 103:104756  
 TITLE: Mechanism of conversion of pyrimido[5,4-e]-1,2,4-triazine-5,7-diones to imidazo[4,5-e]-1,2,4-triazin-6-ones in alkaline media  
 AUTHOR(S): Chernyshev, A. I.; Shorshnev, S. V.; Yakushina, N. I.; Esipov, S. E.  
 CORPORATE SOURCE: Vses. Nauchno-Issled. Inst. Antibiot., Moscow, 113105, USSR  
 SOURCE: Khimiya Geterotsiklicheskikh Soedinenii (1985), (2), 277-8  
 CODEN: KGSSAQ; ISSN: 0453-8234  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 OTHER SOURCE(S): CASREACT 103:104756  
 GI




I



II



III



IV


AB Ring cleavage of reumycin (I; R = H) in alkaline solution of pH .apprx.12 gave the triazinecarboxylate II (R = H) which cyclized at pH >13 to give the imidazolotriazinecarboxylate III (R = H). Decarboxylation of the latter and then oxidation by KMnO4 gave the imidazotriazine IV (R = H). Similarly, fervenulin (I; R = Me) was converted to IV (R = Me).

IT 97943-27-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and decarboxylation of)

RN 97943-27-6 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 1,2,5,6-tetrahydro-5-methyl-6-oxo- (9CI) (CA INDEX NAME)




IT 97943-30-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 97943-30-1 CAPLUS

CN 4aH-Imidazo[4,5-e]-1,2,4-triazine-4a-carboxylic acid, 2,5,6,7-tetrahydro-  
5,7-dimethyl-6-oxo- (9CI) (CA INDEX NAME)



L4 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1977:535265 CAPLUS

DOCUMENT NUMBER: 87:135265

TITLE: Activation and transfer of oxygen. XII. Alkoxy  
adducts derived from 1,3,10-trimethylalloxazinium  
(1,3-dimethylflavinium) cations

AUTHOR(S): Mager, H. I. X.

CORPORATE SOURCE: Biochem. Biophys. Lab., Univ. Technol., Delft, Neth.

SOURCE: Tetrahedron (1977), 33(9), 981-9


CODEN: TETRAB; ISSN: 0040-4020

DOCUMENT TYPE: Journal

LANGUAGE: English

GI





AB Alloxazinium cations with alcs. in the presence of base gave monoalkoxy adducts by nucleophilic attack at C-10a. Dialkoxy adducts were formed by subsequent inter- or intramol. nucleophilic attack at C-4 or C-4a. E.g., I with MeOH gave a 4,10a-dialkoxy adduct, by intermol. attack at C-4, which underwent ring contraction to the hexahydroimidazo[4,5-b]quinoxaline II. I with ethylene glycol gave the 4a,10a-ethylenedioxytetrahydroalloxazine III by intramol. attack at C-4a. Thus, the conclusion of V. Massey and P. Hemmerich (1975) and F. Mueller et al. (1976) that the primary addition site is C-9a is in error. Several rearrangements of II were studied.

IT **64270-53-7P**

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and heterolysis of)


RN 64270-53-7 CAPLUS

CN 1H-Imidazo[4,5-b]quinoxalinium, 9-acetyl-2,4,9,9a-tetrahydro-9a-(methoxycarbonyl)-1,3,4-trimethyl-2-oxo-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 64270-52-6

CMF C16 H19 N4 O4




ONE OR MORE TAUTOMERIC DOUBLE BONDS NOT DISPLAYED IN THE STRUCTURE

CM 2

CRN 14797-73-0

CMF Cl O4



IT 64267-62-5P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation, demethoxycarbonylation, rearrangement, and transesterification of)


RN 64267-62-5 CAPLUS

CN 1H-Imidazo[4,5-b]quinoxalinium, 2,4,9,9a-tetrahydro-9a-(methoxycarbonyl)-  
1,3,4-trimethyl-2-oxo-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 64267-61-4

CMF C14 H17 N4 O3




ONE OR MORE TAUTOMERIC DOUBLE BONDS NOT DISPLAYED IN THE STRUCTURE

CM 2

CRN 14797-73-0

CMF Cl O4



=> log y

COST IN U.S. DOLLARS

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

ENTRY

35.93

SINCE FILE

ENTRY

TOTAL

SESSION

197.47

TOTAL

SESSION

CA SUBSCRIBER PRICE

-5.11

-5.11

STN INTERNATIONAL LOGOFF AT 15:06:04 ON 06 APR 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTAAJP1626

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International\*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 FEB 25 CA/CAPLUS - Russian Agency for Patents and Trademarks  
(ROSPATENT) added to list of core patent offices covered  
NEWS 4 FEB 28 PATDPAFULL - New display fields provide for legal status  
data from INPADOC  
NEWS 5 FEB 28 BABS - Current-awareness alerts (SDIs) available  
NEWS 6 FEB 28 MEDLINE/LMEDLINE reloaded  
NEWS 7 MAR 02 GBFULL: New full-text patent database on STN  
NEWS 8 MAR 03 REGISTRY/ZREGISTRY - Sequence annotations enhanced  
NEWS 9 MAR 03 MEDLINE file segment of TOXCENTER reloaded  
NEWS 10 MAR 22 KOREAPAT now updated monthly; patent information enhanced  
NEWS 11 MAR 22 Original IDE display format returns to REGISTRY/ZREGISTRY  
NEWS 12 MAR 22 PATDPASPC - New patent database available  
NEWS 13 MAR 22 REGISTRY/ZREGISTRY enhanced with experimental property tags  
NEWS 14 APR 04 EPFULL enhanced with additional patent information and new  
fields  
NEWS 15 APR 04 EMBASE - Database reloaded and enhanced

NEWS EXPRESS JANUARY 10 CURRENT WINDOWS VERSION IS V7.01a, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 10 JANUARY 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 15:10:25 ON 06 APR 2005

=> fil reg

COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE

ENTRY

0.42

TOTAL

SESSION

0.42

10/681,924;

5-membered rings as ring "A"

G<sub>2</sub> = N/C



10/681,924  
no art

FILE 'REGISTRY' ENTERED AT 15:11:47 ON 06 APR 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1  
DICTIONARY FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.



```
*****
*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*
*****
```

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10681924i.str



chain nodes :

10 14

ring nodes :

1 2 3 4 5 6 7 8

chain bonds :

5-14 7-10

ring bonds :

1-2 1-5 2-3 3-4 4-5 4-6 5-8 6-7 7-8

exact/norm bonds :

1-2 1-5 2-3 3-4 4-5 4-6 5-14 5-8 6-7 7-8 7-10

G1:O,S

G2:C,N

G3: Cy,Ak

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 10:CLASS 14:CLASS

L1 STRUCTURE UPLOADED

=> d

L1 HAS NO ANSWERS

L1 STR



G1 O,S

G2 C,N

G3 Cy,Ak

Structure attributes must be viewed using STN Express query preparation.

=> s l1 full

FULL SEARCH INITIATED 15:12:10 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 233513 TO ITERATE

100.0% PROCESSED 233513 ITERATIONS

3 ANSWERS

SEARCH TIME: 00.00.06

L2 3 SEA SSS FUL L1

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

161.33

161.75

FILE 'CAPLUS' ENTERED AT 15:12:19 ON 06 APR 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

This file contains CAS Registry Numbers for easy and accurate  
substance identification.

=> s L2

L3 10 L2

=> d L3 1-10 ibib abs hitstr

L3 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1990:157466 CAPLUS

DOCUMENT NUMBER: 112:157466

TITLE: Kinetics of oxidative ammonolysis of copper(2+)  
2,6,8-trioxopurine complex in ammonium buffer

AUTHOR(S): Titskii, G. D.; Miroshnichenko, N. A.

CORPORATE SOURCE: Inst. Phys.-Org. Chem. Coal Chem., Donetsk, USSR

SOURCE: Organic Reactivity (Tartu) (1988), 25(2), 160-9  
CODEN: ORREDZ; ISSN: 0131-8314

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The kinetics of oxidizing conversion of Cu<sup>2+</sup>-2,6,8-trioxypurine complex  
(I) has been studied spectrophotometrically in ammonium buffer (pH  
8.5-10.8) at 25°. The process results from rapid oxidation of I with  
formation of 1-carboxy-2,4,6,8-tetraaza-3,7-dioxo-4-bicyclo[3.3.0]octene  
which is bound into Cu<sup>2+</sup>-NH<sub>3</sub> as complex (II). Slow conversion of (II)  
takes place in two parallel paths.


IT 81129-52-4D, copper ammonium complexes

RL: PRP (Properties)

(intermediacy of, in oxidative ammonolysis of uric acid complex with  
cupric ion, kinetics and mechanism and UV in relation to)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-  
dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1987:523129 CAPLUS

DOCUMENT NUMBER: 107:123129

TITLE: Electrochemical oxidation of 9-β-D-  
ribofuranosyluric acid in basic solution

AUTHOR(S): Tyagi, S. K.; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial  
Electrochemistry (1987), 223(1-2), 119-41  
CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The electrochem. oxidation of 9-β-D-ribofuranosyluric acid (I) was  
studied at the pyrolytic graphite electrode at pH 9.2. The peak Ia  
reaction was a 2e<sup>-</sup>-1H<sup>+</sup> quasi-reversible process giving an unstable  
quinonoid which was rapidly attacked by H<sub>2</sub>O to give 1 and perhaps 2  
tertiary alc. intermediates. In the absence of phosphate, these

intermediates decompose to give a mixture of 5-hydroxyhydantoin-5-carboxamide-3-ribose, ribose and 4-amino-4-carboxy-2,5-diketo-imidazole. In the presence of phosphate, the putative tertiary alc. intermediate underwent a ring contraction to an electrooxidizable bicyclic carboxylic acid which decomposed to an allantoin riboside. The carboxylic acid was partially responsible for the 2nd, more pos. oxidation peak of I (peak IIa) and in a 2e--1H+ reaction gave 1-hydroxy-2,4,6,8-tetraaza-3,7-dioxo-5-ene-bicyclo(3.3.0)-octane-4-ribose. A stable, peak Ia product also contributed to the peak IIa reaction. This species was a phosphorylated derivative of the bicyclic carboxylic acid. At pH 9.2, in the presence of phosphate, type VIII peroxidase was capable of oxidizing I in a chemical reaction equivalent to the peak Ia electrochem. process.


IT **85303-15-7P**

RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, from rearrangement of tertiary alc.)

RN 85303-15-7 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-dioxo-1-β-D-ribofuranosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L3 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1986:157924 CAPLUS

DOCUMENT NUMBER: 104:157924

TITLE: Electrochemical oxidation of 2,6-diamino-8-purinol

AUTHOR(S): Astwood, D.; D'Amico, C. N.; Lippincott, T.;  
Brajer-Toth, A.

CORPORATE SOURCE: Dep. Chem., Univ. Maine, Orono, ME, 04469, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial  
Electrochemistry (1986), 198(2), 283-302  
CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Electrochem., spectroelectrochem. and gas chromatog./mass spectrometry were used to determine the mechanism of oxidation of 2,6-diamino-8-purinol [72308-52-2] in phosphate buffer solns. Exptl. evidence indicates that the process is a 2 e-, 2 H+ reaction in which 2,6-diamino-8-purinol is oxidized to form a diimine. The diimine is not stable and decomp. in a series of hydrolysis reactions to the final products 5-hydroxyhydantoin-5-carboxamide [36597-25-8] and 3-carbohydroxy-2,4,6,8-tetraza-3,7-dioxo-4-ene-bicyclo[3.3.0]octane. Results are presented which show that the 2 e- oxidation of 2,6-diamino-8-purinol proceeds in two 1 e-1 steps and the evidence for the formation of a coupling product is discussed. A mechanism is proposed to explain the observed results.

IT **81129-52-4P**


RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in electrochem. oxidation of diaminopurinol in phosphate buffered solution)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-



dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1984:93328 CAPLUS

DOCUMENT NUMBER: 100:93328

TITLE: Electrochemical and enzymatic oxidation of  
2,6-diaminopurine

AUTHOR(S): Astwood, D.; Lippincott, T.; Deysher, M.; D'Amico, C.;  
Szurley, E.; Brajter-Toth, A.

CORPORATE SOURCE: Dep. Chem., Univ. Maine, Orono, ME, 04469, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial  
Electrochemistry (1983), 159(2), 295-312  
CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Electrochem. spectroelectrochem., gas chromatog./mass spectrometry (GC/MS) were used to study the mechanism of electrochem. oxidation of 2,6-diaminopurine [1904-98-9] in aqueous solns. Exptl. evidence indicates that the oxidation pathway depends on potential. At less pos. potentials 2,6-diaminopurine is initially oxidized in a 2 e<sup>-</sup>, 2 H<sup>+</sup> reaction to 2,6-diamino-8-purinol [72308-52-2] which is immediately further oxidized to the 2,6-diaminopurinol-diimine in a 2 e<sup>-</sup>, 2 H<sup>+</sup> reaction. At more pos. potentials 4 e and 4 H<sup>+</sup> are exchanged to form the diimine. The diimine is not stable and decomp. in a series of hydrolysis reactions to the final products 5-hydroxyhydantoin-5-carboxamide [36597-25-8] and 1-carbohydroxy-2,4,6,8-tetraaza-3,7-dioxo-4-ene-bicyclo[3.3.0]octane [81129-52-4]. The products were identified by GC/MS. A mechanism is proposed to explain the observed results. The significance of this mechanism with respect to the enzymic oxidation and deamination of 2,6-diaminopurine is discussed.


IT 81129-52-4P

RL: FORM (Formation, nonpreparative); PREP (Preparation)

(formation of, in electrochem. oxidation of diaminopurine in aqueous solns.)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1983:620927 CAPLUS

DOCUMENT NUMBER: 99:220927

TITLE: Electrochemical oxidation of 7,9-dimethyluric acid in  
acid solution

AUTHOR(S): Chen, Tsuyu Raymond; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial

DOCUMENT TYPE: Journal  
LANGUAGE: English


AB The electrochem. oxidation of 7,9-dimethyluric acid [19039-41-9] at pH 3-5 was studied by using a pyrolytic graphite electrode. Two voltammetric oxidation peaks (Ia and IIa) were observed. Peak Ia is a  $2e^- - 1H^+$  reaction yielding a very reactive quinoid cation that, in a reaction catalyzed by  $H_2PO_4^-$ , is hydrated to an UV-absorbing tertiary alc. intermediate. This intermediate can undergo ring contraction to give 1-carboxy-2,4-dimethyl-2,4,6,8-tetraaza-3,7-dioxo-5-ene-bicyclo[3.3.0]octane (I) [85303-22-6] or react with  $H_2PO_4^-$  and 7,9-dimethyluric acid to give a relatively long-lived complex. The I is hydrated and decomp. to 1,3-dimethylallantoin [32282-45-4]. The tertiary alc.- $H_2PO_4^-$ -7,9-dimethyluric acid complex very slowly decomp. regenerating 7,9-dimethyluric acid and, ultimately, 1,3-dimethyl-5-hydroxyhydantoin-5-carboxamide [87897-69-6]. Peak IIa is a minor process and is due to  $2e^-$  oxidation of the anion of I in a Kolbe-type reaction giving 1-hydroxy-2,4-dimethyl-2,4,6,8-tetraaza-3,7-dioxo-5-ene-bicyclo[3.3.0]octane [85303-23-7].

IT 85303-22-6P

RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in electrochem. oxidation of dimethyluric acid in acid phosphate buffer solution)

RN 85303-22-6 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-1,3-dimethyl-2,5-dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1983:169134 CAPLUS

DOCUMENT NUMBER: 98:169134

TITLE: The electrochemical and peroxidase-catalyzed redox chemistry of 9- $\beta$ -D-ribofuranosyluric acid

AUTHOR(S): Goyal, R. N.; Brajter-Toth, Anna; Besca, Joseph S.; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial Electrochemistry (1983), 144(1-2), 163-90  
CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The electrochem. oxidation of 9- $\beta$ -D-ribofuranosyluric acid (I) [21082-30-4] at a pyrolytic graphite electrode proceeds via an initial  $2e^-$  reaction to give an unstable quinonoid intermediate. Nucleophilic attack by water of this intermediate leads to isomeric tertiary alc. intermediates. The latter were characterized by electrochem. reduction to a dihydro product which readily dehydrates to regenerate I. The tertiary alc. intermediates can undergo a ring-opening reaction at pH  $\geq 6$  to give a spectrally distinct and reversibly reducible pyrimidine derivative(s) which slowly decompose to give alloxan or alloxanic acid and urea riboside. Alternatively, the tertiary alc. can further hydrate to yield, ultimately,

5-hydroxyhydrantoin-5-carboxamide-3-riboside [85303-13-5] or undergo a ring contraction reaction and hydrolysis leading to allantoin riboside [85303-14-6]. The peroxidase-catalyzed oxidation appears to follow essentially the same chemical pathway.


IT **85303-15-7P**

RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in electrochem. oxidation of ribofuranosyluric acid)

RN 85303-15-7 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-dioxo-1-β-D-ribofuranosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L3 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1983:151660 CAPLUS

DOCUMENT NUMBER: 98:151660

TITLE: The second voltammetric oxidation peak of  
7,9-dimethyluric acid

AUTHOR(S): Chen, Tsuyu Raymond; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial  
Electrochemistry (1983), 144(1-2), 191-206  
CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Voltammetry of 7,9-dimethyluric acid [19039-41-9] in aqueous solution at a pyrolytic graphite electrode shows that 2 oxidation peaks are formed. Peak Ia is a 2-e electrooxidn. giving a very unstable quinonoid cation. This rapidly hydrates in a phosphate-catalyzed reaction giving a tertiary alc., which in turn undergoes a ring contraction to 1-carbohydroxy-2,4-dimethyl-2,4,6,8-tetraaza-3,7-dioxo-5-enebicyclo[3.3.0]octane [85303-22-6]. The latter is again unstable and hydrates and decarboxylates, giving 1,3-dimethylallantoin [32282-45-4] as the final product. Peak IIa is due to a Kolbe-type 2-e electrooxidn. of 1-carbohydroxy-2,4-dimethyl-2,4,6,8-tetraaza-3,7-dioxo-5-enebicyclo[3.3.0]octane to give a very reactive carbonium ion and CO<sub>2</sub>. The former species is rapidly hydrated, giving 1-hydroxy-2,4-dimethyl-2,4,6,8-tetraaza-3,7-dioxo-5-enebicyclo[3.3.0]octane [85303-23-7], a new compound which has been isolated and characterized by spectral, chemical and electrochem. methods.

IT **85303-22-6**

RL: PRP (Properties)  
(electrochem. oxidation of electrogenerated intermediate of, in  
dimethyluric acid electrochem. oxidation)

RN 85303-22-6 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-1,3-dimethyl-2,5-dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1982:571170 CAPLUS

DOCUMENT NUMBER: 97:171170

TITLE: Spectroelectrochemical evidence for imine-alcohol intermediate formed upon electrochemical oxidation of uric acid

AUTHOR(S): Goyal, R. N.; Nguyen, N. T.; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Bioelectrochemistry and Bioenergetics (1982), 9(3), 273-85

CODEN: BEBEBP; ISSN: 0302-4598

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Electrochem. oxidation of uric acid [69-93-2] in phosphate-containing supporting

electrolytes at pH 3-9 at a reticulated vitreous C electrode in a thin-layer spectroelectrochem. cell leads to formation of UV-absorbing intermediate species. Electrochem. reduction of the intermediate-containing solution

leads to the partial regeneration of uric acid. This behavior provides compelling evidence that an imine-alc. is 1 of the UV-absorbing intermediate species because only this compound may be expected to be reduced to a species which can regenerate uric acid.


IT 81129-52-4

RL: PRP (Properties)

(intermediate, in uric acid electrochem. oxidation)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 9 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1982:467958 CAPLUS

DOCUMENT NUMBER: 97:67958

TITLE: A comparison of the peroxidase-catalyzed and electrochemical oxidation of uric acid

AUTHOR(S): Goyal, R. N.; Brajter-Toth, Anna; Dryhurst, Glenn; Nguyen, N. T.

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Bioelectrochemistry and Bioenergetics (1982), 9(1), 39-60

CODEN: BEBEBP; ISSN: 0302-4598

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The oxidation of uric acid by H<sub>2</sub>O<sub>2</sub> in the presence of type VIII peroxidase was studied between pH 5.2 and 8. Intermediates generated in the reaction were characterized in terms of their UV spectra and kinetics of decay. In addition ≥1 UV-absorbing intermediate was trapped, converted to its trimethylsilyl derivative, and identified by gas chromatog.-mass spectrometry. This intermediate is 1-carbohydroxy-2,4,6,8-tetraza-3,7-dioxo-4-ene-bicyclo-(3.3.0)-octane. At pH ≥7 the product is allantoin, whereas at lower pH 5-hydroxyhydantoin-5-carboxamide is also formed as a major product. The intermediates and products formed and spectral and kinetic measurements observed during and after peroxidase-catalyzed oxidation of uric acid are virtually identical to those noted upon electrochem. oxidation. Thus, the mechanisms of electrochem. and enzymic oxidation of uric acid are, in a chemical sense, identical.


IT **81129-52-4**

RL: BIOL (Biological study)

(as urate peroxidase-catalyzed oxidation intermediate)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-dioxo- (9CI) (CA INDEX NAME)



L3 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1982:151261 CAPLUS

DOCUMENT NUMBER: 96:151261

TITLE: Further insights into the electrochemical oxidation of uric acid

AUTHOR(S): Goyal, R. N.; Brajter-Toth, Anna; Dryhurst, Glenn

CORPORATE SOURCE: Dep. Chem., Univ. Oklahoma, Norman, OK, 73019, USA

SOURCE: Journal of Electroanalytical Chemistry and Interfacial Electrochemistry (1982), 131, 181-202

CODEN: JEIEBC; ISSN: 0022-0728

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The electrochem. oxidation of uric acid [69-93-2] was studied between pH 1.5 and 9.5 in phosphate buffers using thin-layer spectroelectrochem. to generate and study UV-absorbing intermediates. It was concluded, on the basis of this and preceding studies, that uric acid is 1st oxidized in a 2e--2H+ reaction to a very unstable quinonoid diimine (half-life ≤ 22 ms). At pH ≥ 6 the anion of the latter species is attacked by water to give an anionic imine-alc. that undergoes a ring contraction reaction to give 1-carbohydroxy-2,4,6,8-tetraaza-3,7-dioxo-4-enebicyclo[3.3.0]octane [81129-52-4]. This then decomp. to allantoin [97-59-6]. At pH 3-5.6 a neutral quinonoid diimine is generated upon 2e--2H+ oxidation of uric acid. In high-phosphate buffers H<sub>2</sub>PO<sub>4</sub><sup>-</sup> attacks the diimine, whereas in low-phosphate buffers solvent (H<sub>2</sub>O) attacks the diimine. In high-phosphate buffers anal. of absorbance vs. time curves obtained following oxidation of uric acid in a thin-layer cell allows 3 intermediate species to be inferred. In low-phosphate buffers only 2 intermediates may be inferred. Mechanisms are advanced to rationalize these observations and to account for the end products formed, i.e. allantoin, 5-hydroxyhydantoin-5-carboxamide [36597-25-8] and, at pH 3, alloxan [50-71-5].


IT **81129-52-4P**

RL: FORM (Formation, nonpreparative); PREP (Preparation)

(formation of, in electrochem. oxidation of uric acid in  
phosphate-buffered solution)

RN 81129-52-4 CAPLUS

CN Imidazo[4,5-d]imidazole-3a(1H)-carboxylic acid, 2,3,4,5-tetrahydro-2,5-  
dioxo- (9CI) (CA INDEX NAME)



=> fil reg

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
51.20	212.95

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-7.30	-7.30

CA SUBSCRIBER PRICE

FILE 'REGISTRY' ENTERED AT 15:14:31 ON 06 APR 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1  
DICTIONARY FILE UPDATES: 5 APR 2005 HIGHEST RN 847968-12-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

\*\*\*\*\*  
\*  
\* The CA roles and document type information have been removed from \*  
\* the IDE default display format and the ED field has been added, \*  
\* effective March 20, 2005. A new display format, IDERL, is now \*  
\* available and contains the CA role and document type information. \*  
\*  
\*\*\*\*\*

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10681924j.str